

## Note

# Combinatorial Interpretation and Operator Calculus of Lommel Polynomials

P. FEINSILVER AND J. MCSORLEY

*Department of Mathematics, Southern Illinois University, Carbondale, Illinois 62901*

AND

R. SCHOTT



CORE

Provided by Elsevier - Publisher Connector

Received July 28, 1995

In this paper we present interpretations of Lommel polynomials and their derivatives. A combinatorial interpretation uses matchings in graphs. This gives an interpretation for the derivatives as well. Then Lommel polynomials are considered from the point of view of operator calculus. A step-3 nilpotent Lie algebra and finite-difference operators arise in the analysis. © 1996 Academic Press, Inc.

## I. INTRODUCTION

Interpretations of orthogonal polynomials in terms of combinatorial models has received a lot of attention over the last decade. Dulucq and Favreau [3] recently presented a combinatorial model for Bessel polynomials. A general combinatorial theory for orthogonal polynomials has been developed in the work of Viennot [12], de Médicis and Viennot [2], and in the theory of species, formalized by Bergeron [1], Joyal [6, 7], Labelle [8], and Leroux [9]. An analytical study of the zeros of Lommel polynomials may be found in [5]. The basics of the operator calculus approach are in [4].

Lommel polynomials arise in the study of Bessel functions as the linearization coefficients expressing  $J_{v+n}$  in terms of  $J_v$  and  $J_{v-1}$ ; cf. Watson [13]. They may be given explicitly in the form

$$R_n(\xi, v) = \sum_{k=0}^{\lfloor n/2 \rfloor} \binom{n-k}{k} (-1)^k \frac{\Gamma(v+n-k)}{\Gamma(v+k)} (2/\xi)^{n-2k}.$$

Changing variables to  $\phi_n(x, \varepsilon) = R_n(-2/\varepsilon, -x/\varepsilon)$ , with  $\varepsilon$  understood as a parameter, we have the recurrence

$$x\phi_n = \phi_{n+1} + \varepsilon n\phi_n + \phi_{n-1} \quad (1.1)$$

with initial conditions  $\phi_{-1} = 0$ ,  $\phi_0 = 1$ . Thus, these are a family of orthogonal polynomials. They have the form

$$\phi_n(x, \varepsilon) = \sum_k \binom{n-k}{k} (-1)^k (x - k\varepsilon) \cdots (x - (n-k-1)\varepsilon). \quad (1.2)$$

In this paper we present some interpretations of Lommel polynomials. We proceed from a general approach and then specialize to the case of Lommel polynomials. For example, note that with  $\varepsilon = 0$  in Eq. (1.1), we have the recurrence for Chebyshev polynomials. In fact, with  $\varepsilon = 0$ , the polynomials  $\phi_n$  become the Chebyshev polynomials of the second kind.

## II. BASIC CONSTRUCTION: MATCHINGS

Let  $G$  be a simple graph on  $n$  vertices with vertex labels 1 to  $n$  and weight  $w_i$  on vertex  $i$ ,  $1 \leq i \leq n$ . A *matching*,  $M$ , of  $G$  is a set of disjoint edges pairwise having no vertex in common. If a vertex  $i$  is incident to an edge of  $M$ , we write  $i \in M$ ; otherwise  $i \notin M$ . We define the weight of  $M$ ,  $W_G(M)$ , to be

$$W_G(M) = \prod_{i \notin M} w_i.$$

If  $G$  has an even number of vertices and  $M$  is a perfect matching of  $G$ , in other words, if all vertices of  $G$  lie in  $M$ , then  $W_G(M)$  is defined to equal 1. Note that every graph has the empty matching, containing no edges.

For the rest of this work, we will take  $G$  to be  $P_n$ , the path on  $n$  vertices, running from left to right. We denote  $W_G(M)$ , then, by  $W_n(M)$ , the weight of the matching  $M$  of  $P_n$ .

Define

$$\mathcal{P}_n = \sum_M (-1)^{|M|} W_n(M)$$

with  $|M|$  the number of edges in  $M$ , summing over all matchings  $M$  of  $P_n$ .

2.1. PROPOSITION.  $\mathcal{P}_n$  satisfies the recurrence

$$\mathcal{P}_{n+1} = w_{n+1} \mathcal{P}_n - \mathcal{P}_{n-1}$$

*Proof.* Let  $M$  be a matching of  $\mathcal{P}_{n+1}$ . Either  $M$  does not contain the edge  $(n, n+1)$  or it does. Every case where the edge  $(n, n+1)$  is not in  $M$  corresponds to some matching on  $n$  vertices with the additional factor of  $w_{n+1}$  since  $n+1$  is in none of them. On the other hand, in all matchings including the edge  $(n, n+1)$ , removing it leaves a matching on  $n-1$  vertices with the removal of the edge contributing a minus sign. ■

We define  $\mathcal{P}_{-1}=0$ ,  $\mathcal{P}_0=1$ . From the proposition, this gives  $\mathcal{P}_1=w_1$ , which agrees with the scheme, since the only matching of  $P_1$ , a single vertex, is the empty matching, with weight  $w_1$ . Similarly, we have from the proposition that  $\mathcal{P}_2=w_1w_2-1$ . The graph  $P_2$  has two matchings—the empty matching with weight  $w_1w_2$  and the matching which contains the single edge  $(1, 2)$  of weight 1.

Now recall that a three-term recurrence of the form

$$x\phi_n = \phi_{n+1} + b_n\phi_n + c_n\phi_{n-1}$$

with  $c_n \geq 0$  yields a sequence of orthogonal polynomials. In Proposition 2.1, we can introduce a variable  $x$  either additively or multiplicatively into the weights. I.e., consider

$$(x - u_{n+1})\mathcal{P}_n = \mathcal{P}_{n+1} + \mathcal{P}_{n-1}$$

which means that  $x$  as the eigenvalues of the corresponding tridiagonal matrices are the zeros of the polynomials  $\mathcal{P}_n$ . Or we can write  $w_{n+1} = xu_{n+1}$  with the recurrence taking the form

$$xu_{n+1}\mathcal{P}_n = \mathcal{P}_{n+1} + \mathcal{P}_{n-1},$$

for example, with constant  $u_{n+1}=2$ , we have the recurrence for the Chebyshev polynomials.

## 2.1. Lommel Polynomials and Their Derivatives

The recurrence (1.1) corresponds to the weights  $w_i = x - (i-1)\varepsilon$ . For fixed  $\varepsilon$ , we treat  $\phi_n$  as a function of one variable,  $x$ , and denote it by  $R_n$ . Here are some explicit expressions:

$$R_1 = x, \quad R_3 = x^3 - 3\varepsilon x^2 + 2\varepsilon^2 x - 2x + 2\varepsilon,$$

$$R_2 = x^2 - \varepsilon x - 1, \quad R_4 = x^4 - 6\varepsilon x^3 + 11\varepsilon^2 x^2 - 6\varepsilon^3 x - 3x^2 + 9\varepsilon x - 6\varepsilon^2 + 1.$$

Now, let  $G$  be a simple graph with  $n$  vertices labelled 1 to  $n$  having weight  $x - (i-1)\varepsilon$  on vertex  $i$ . Define a  $k$ -extended matching of  $G$  as a set  $\{v_1, \dots, v_k, M\}$  of  $k$  vertices of  $G$ , together with a matching  $M$  such that  $v_j \notin M$  for  $1 \leq j \leq k$ . I.e., no vertex  $v_j$  is incident to any edge in  $M$ .

Denoting the  $k$ -extended matching  $\{v_1, \dots, v_k, M\}$  by  $E_M$ ,  $M$  is called the matching of  $E_M$ . For a vertex  $v \in G$ , we write  $v \in E_M$  if either  $v = v_j$  for some  $j$ ,  $1 \leq j \leq k$ , or  $v \in M$ . Let  $|E_M| = |M|$  denote the number of edges in the matching of  $E_M$ .

The weight of  $E_M$ ,  $W_G(E_M)$  is given by

$$W_G(E_M) = \prod_{i \notin E_M} (x - (i-1)\varepsilon).$$

As above, if every vertex of  $G$  lies in  $E_M$ , then  $W_G(E_M) = 1$ . We take  $G = P_n$ , the path, and denote  $W_G(E_M)$  by  $W_n(E_M)$ .

### 2.1.1. Derivatives of Lommel Polynomials

For a combinatorial interpretation of the  $k$ th derivative  $R_n^{(k)}$ , start from the contribution of the matching  $M$ ,  $W_n(M) = \prod_{i \notin M} (x - (i-1)\varepsilon)$ . Consider  $W_n(M)$  as a function of  $x$ . Since the derivative of each factor in  $W_n(M)$  is 1, for the  $k$ th derivative we have, the summation taken over all  $k$ -subsets of vertices  $\{v_{i_1}, \dots, v_{i_k}\}$  of  $G$  such that no vertex  $v_{i_j}$  is in  $M$ ,

$$\begin{aligned} W_n^{(k)}(M) &= \sum_{\{i_1, \dots, i_k\}} \frac{k! W_n(M)}{\prod_j (x - (i_j-1)\varepsilon)} \\ &= k! \sum_{\substack{\{i_1, \dots, i_k\} \\ v_{i_j} \notin M}} W_n(\{i_1, \dots, i_k, M\}) \\ &= k! \sum_{E_M} W_n(E_M); \end{aligned}$$

this last is summation taken over all  $k$ -extended matchings  $E_M$  with matching  $M$ . Hence

**2.1.1.1. PROPOSITION.** The  $k$ th derivative of the Lommel polynomial  $R_n$  is given by

$$R_n^{(k)} = k! \sum_E (-1)^{|E|} W_n(E),$$

where the summation is over all  $k$ -extended matchings of  $P_n$ , with  $|E|$  denoting the number of edges in the matching of  $E$ .

Note that with  $k=0$  we recover the original case of  $R_n$ , considering a matching as a 0-extended matching.

**EXAMPLE.** Consider the second derivative of  $R_4$ . We have

$$\begin{aligned} R_4 &= x^4 - 6\varepsilon x^3 + 11\varepsilon^2 x^2 - 6\varepsilon^3 x - 3x^2 + 9\varepsilon x - 6\varepsilon^2 + 1 \\ \frac{1}{2} R_4'' &= 6x^2 - 18\varepsilon x + 11\varepsilon^2 - 3. \end{aligned}$$

The 2-extended matchings with corresponding weights are given in the following table:

2-ext. match.	Weight
$\{1, 2, \emptyset\}$	$(x - 2\varepsilon)(x - 3\varepsilon)$
$\{1, 3, \emptyset\}$	$(x - \varepsilon)(x - 3\varepsilon)$
$\{1, 4, \emptyset\}$	$(x - \varepsilon)(x - 2\varepsilon)$
$\{2, 3, \emptyset\}$	$x(x - 3\varepsilon)$
$\{2, 4, \emptyset\}$	$x(x - 2\varepsilon)$
$\{3, 4, \emptyset\}$	$x(x - \varepsilon)$
$\{3, 4, (1, 2)\}$	1
$\{1, 4, (2, 3)\}$	1
$\{1, 2, (3, 4)\}$	1

### III. LOMMEL POLYNOMIALS AND FINITE-DIFFERENCE CALCULUS

Here we show how to write Lommel polynomials in terms of a recurrence with operator coefficients. Recall that the solution to the recurrence

$$f_{n+1} = af_n + bf_{n-1}$$

with initial conditions  $f_{-1} = 0, f_0 = 1$ , is given by

$$f_n = \sum_k \binom{n-k}{k} a^{n-2k} b^k.$$

(In other terms, we can express the solution to

$$f_{n+1} = af_n - bf_{n-1}$$

with the same initial conditions, in terms of Chebyshev polynomials of the second kind:  $f_n = b^{n/2} U_n(a/(2\sqrt{b}))$ .) This formula holds for  $a$  and  $b$  operators, e.g., matrices, with  $f_0 = I$ , the identity, as long as  $a$  and  $b$  commute. If they do not commute, we apply them on different sides.

**3.1. PROPOSITION.** *For operators  $a$  and  $b$ , the solution to the recurrence*

$$f_{n+1} = f_n a + b f_{n-1}$$

*with initial conditions  $f_{-1} = 0, f_0 = I$ , is given by*

$$f_n = \sum_k \binom{n-k}{k} b^k a^{n-2k},$$

*and, similarly, with  $a$  acting on the left and  $b$  on the right.*

For Lommel polynomials, introduce the shift operator  $T_\varepsilon$  acting on functions  $f$  by

$$T_\varepsilon f(x) = f(x - \varepsilon).$$

We denote the operator of multiplication by  $x$  by  $X$ . Using the relation

$$(XT_\varepsilon)^n 1 = x(x - \varepsilon)(x - 2\varepsilon) \cdots (x - (n - 1)\varepsilon),$$

where 1 denotes the constant function 1, we have from Eq. (1.2),

$$R_n(x) = \sum_k \binom{n-k}{k} (-T_\varepsilon)^k (XT_\varepsilon)^{n-2k} 1. \quad (3.1)$$

Comparing with Proposition 3.1, we see the following.

3.2. PROPOSITION. *Define operators  $F_n$  by the recurrence*

$$F_{n+1} = F_n XT_\varepsilon - T_\varepsilon F_{n-1}$$

*with  $F_{-1} = 0$ ,  $F_0 = I$ . Then the Lommel polynomials are given by*

$$R_n(x) = F_n 1.$$

*For example,*

$$F_1 = XT_\varepsilon, \quad F_2 = (XT_\varepsilon)^2 - T_\varepsilon, \quad F_3 = (XT_\varepsilon)^3 - 2T_\varepsilon XT_\varepsilon,$$

*etc.*

Another approach to expressions of the form

$$\sum_k \binom{n-k}{k} a^{n-2k} b^k$$

involves the nilpotent Lie algebra generated by the operators  $D^2 = (d/dx)^2$  and  $X$ . The commutator  $[D^2, X] = D^2 X - X D^2 = 2D$ , while  $[D, X] = 1$ . Thus,  $\{D^2, D, X, 1\}$  form the basis for a nilpotent Lie algebra of step 3; i.e., all commutators of length greater than 3 vanish. Now, we have the following.

3.3. PROPOSITION. *Let  $f_n(x) = \sum_k \binom{n-k}{k} b^k x^{n-2k}$ . Then*

$$f_n(x) = {}_0F_1 \left( \begin{matrix} - \\ -n \end{matrix} \middle| -bD^2 \right) x^n.$$

*Proof.* Expanding the  ${}_0F_1$  function gives

$$\sum_k \frac{(-bD^2)^k}{(-n)_k k!} x^n = \sum_k \frac{(n-k)!}{n! k!} b^k D^{2k} x^n$$

from which the result is clear. ■

To see the connection with Lommel polynomials, first we review the basic operator calculus needed. Consider the formal series in one variable

$$V(z) = \sum_{n=0}^{\infty} a_n z^n.$$

This is the *symbol* of the generalized differential operator  $V(D)$ , which acts on polynomials in the variable  $x$ . This satisfies

$$[V(D), X] = V'(D). \quad (3.2)$$

$V'(z)$  denotes the derivative of the series  $V(z)$ . We assume that  $a_0=0$ ,  $a_1=1$ . Thus,  $V'$  has a formal multiplicative inverse  $1/V'(z)$ , which we denote by  $W(z)$ . From Eq. (3.2), we see that, defining  $\xi = XW(D)$ , we have

$$[V(D), \xi] = 1,$$

from which the usual rules of polynomial calculus follow, such as

$$V\xi^n 1 = n\xi^{n-1} 1.$$

The shift operator  $T_\varepsilon$  has symbol  $e^{-\varepsilon z}$ . Thus, for the operator  $\xi = XT_\varepsilon$ , we have the corresponding operator  $V(D)$  with symbol

$$V(z) = \frac{1}{\varepsilon} (e^{\varepsilon z} - 1) \quad (3.3)$$

which is the (forward) finite-difference operator with step size  $\varepsilon$ . Next, define the *finte-difference Laplacian* with the symbol

$$\Delta_\varepsilon(z) = \frac{1}{\varepsilon^2} (e^{\varepsilon z} + e^{-\varepsilon z} - 2).$$

Now we have the following.

3.4. Proposition. *The Lommel polynomials  $R_n(x)$  satisfy*

$$R_n(x) = {}_0F_1 \left( \begin{matrix} - \\ -n \end{matrix} \middle| \Delta_\varepsilon \right) \xi^n 1$$

with  $\xi = XT_\varepsilon$ .

*Proof.* Write Eq. (3.1) in the form

$$R_n(x) = \sum_k \binom{n-k}{k} (-T_\varepsilon)^k \xi^{n-2k} 1.$$

Then, as in Proposition 3.3,

$$R_n(x) = {}_0F_1 \left( \begin{matrix} - \\ -n \end{matrix} \middle| T_\varepsilon V(D)^2 \right) \xi^n 1$$

with  $V(D)$  the difference operator in Eq. (3.3). Now calculating with symbols, we see that

$$T_\varepsilon(z) V(z)^2 = \frac{1}{\varepsilon} (e^{\varepsilon z} - 1)(1 - e^{-\varepsilon z}) = \Delta_\varepsilon(z)$$

and, hence, the result. ■

#### IV. CONCLUDING REMARKS

It would be interesting to consider the combinatorial approach for other families of orthogonal polynomials as indicated in Section II. In [4], the function  ${}_0F_1$  arises naturally in the  $\mathfrak{sl}(2)$  calculus yielding eigenfunctions of the radial Laplacian in Euclidean space. Here, we see that the Lommel polynomials correspond to the finite-difference Laplacian. It appears that the Lommel polynomials play a natural role in harmonic analysis on a lattice and merit further study in this context.

#### ACKNOWLEDGMENT

This work was carried out with the help of NATO Collaborative Research Grant 931395.

#### REFERENCES

1. F. BERGERON, A combinatorial outlook on symmetric functions, *J. Combin. Theory Ser. A* **50**, No. 2 (1989), 226–234.
2. A. DE MÉDICIS AND X. G. VIENNOT, Moments des  $q$ -Polynômes de Laguerre et la Bijection de Foata-Zeiberger, *Adv. in Appl. Math.* **15** (1994), 262–304.
3. S. DULUCQ AND L. FAVREAU, Un modèle combinatoire pour les polynômes de Bessel, in “Séminaire Lotharingien de Combinatoire, 1990,” Université de Strasbourg, 1991.
4. P. FEINSILVER AND R. SCHOTT, “Algebraic Structures and Operator Calculus,” Vol. 1, Representations and Probability Theory, Kluwer, Amsterdam, 1993.



5. P. FEINSILVER AND R. SCHOTT, On Bessel functions and rate of convergence of zeros of Lommel polynomials, *Math. Comp.* **59** (1992), 199, 153–156.
6. A. JOYAL, Une théorie combinatoire des séries formelles, *Adv. in Math.* **42** (1981), 1–82.
7. A. JOYAL, Foncteurs analytiques et espèces de structures, in “Proceedings, Colloque de Comb. Enumer. UQAM, 1985,” Lectures Notes in Math., Vol. 1234, pp. 126–159, Springer–Verlag, New York/Berlin, 1986.
8. G. LABELLE, Une nouvelle démonstration combinatoire des formules d’inversion de Lagrange, *Adv. in Math.* **42** (1981), 217–247.
9. P. LEROUX AND V. STREHL, Jacobi polynomials: Combinatorics of the basic identities, *Discrete Math.* **57** (1985), 167–187.
10. G. C. ROTA (Ed.), “Finite Operator Calculus,” Academic Press, New York/London, 1975.
11. G. SZÉGO, “Orthogonal Polynomials,” AMS Colloq. Publ., *Amer. Math. Soc.*, Providence, RI, 1975.
12. X. G. VIENNOT, Une théorie combinatoire des polynômes orthogonaux généraux, Notes de cours, Université du Québec à Montréal, 1983.
13. G. N. WATSON, “A Treatise on the Theory of Bessel Functions,” Cambridge Univ. Press, Cambridge, 1980.